

Welcome to NEXLETOL

A once-daily, nonstatin pill for lowering your bad cholesterol (LDL-C)



What is NEXLETOL?

NEXLETOL is a once-daily, nonstatin pill for adults. It is used to help lower your bad cholesterol when you add it to a healthy diet and other lipid-lowering medicines.

NEXLETOL is for adults who have:

- heterozygous familial hypercholesterolemia (an inherited condition that causes high levels of bad cholesterol) and/or
- known cardiovascular disease due to high levels of bad cholesterol

It is not known if NEXLETOL can decrease problems related to high cholesterol, such as heart attacks or stroke.

The active ingredient in NEXLETOL is bempedoic acid. It works by lowering the amount of cholesterol your liver makes. This eventually leads to removal of bad cholesterol from your blood.

For further information about NEXLETOL, [see here](#) for the full Prescribing Information.

The use of NEXLETOL has not been studied in children under 18 years of age.

What are the possible side effects of NEXLETOL?

The most common side effects include:

- Upper respiratory tract infection
- Muscle spasms
- Hyperuricemia
- Back pain
- Abdominal pain or discomfort

Other serious side effects may include:

- Increased levels of uric acid in your blood, which may lead to gout
- Serious tendon problems

For signs and symptoms of these serious side effects, please see "What is the most important safety information I should know about NEXLETOL?" on the following page.

These are not all the possible side effects of NEXLETOL.

You may not notice some of these side effects. Your doctor may need to do a lab test in order to find out if you are experiencing them.

Contact your doctor if you experience or require more information about side effects. You may also report side effects to the Food and Drug Administration at 1-800-FDA-1088.

How should I take NEXLETOL?

Always take this medicine exactly as your doctor prescribes.

- Take 1 pill, once daily, by mouth. **You may take it with or without food**
- If you take more NEXLETOL than you should, call poison control center at 1-800-222-1222 or go to the nearest hospital emergency room right away

What if I am also taking other medicines?

Tell your doctor about all the medicines you take. These include prescription and over-the-counter medicines, vitamins, and herbal supplements.

It's important to tell your doctor if you are taking simvastatin or pravastatin (other medicines for lowering cholesterol) daily.

Taking NEXLETOL with over 20 mg simvastatin daily or over 40 mg pravastatin daily could increase your chance of getting muscle side effects. This means your doctor may need to lower your dose of simvastatin or pravastatin.

What if I miss a dose of NEXLETOL?

Do not take more than 1 dose per day.

If you miss a dose, talk to your doctor.

How should I store NEXLETOL?

- Store NEXLETOL in its original package at room temperature
- Keep NEXLETOL and all medicines out of the reach of children

Am I eligible for co-pay assistance?

- See if you are eligible for co-pay assistance and savings at NexCopay.com



LDL-C=low-density lipoprotein cholesterol.

Stay in the know about NEXLETOL!



Call our information line at
1-833-NEXLETOL (1-833-639-5386)



or visit NEXLETOL.com and sign up to receive more information.

Please see Important Safety Information on the following page and full Prescribing Information.

What is NEXLETOL?

NEXLETOL is a prescription medicine used, along with diet and other lipid-lowering medicines, in the treatment of adults who need additional lowering of “bad” cholesterol (LDL-C) and have:

- heterozygous familial hypercholesterolemia (HeFH), an inherited condition that causes high levels of LDL-C, and/or
- known cardiovascular disease

It is not known if NEXLETOL can decrease problems related to high cholesterol, such as heart attacks or stroke.

IMPORTANT SAFETY INFORMATION

What should I tell my doctor before taking NEXLETOL?

Tell your doctor if you:

- have, or have ever had, gout
- have, or have ever had, tendon problems
- are pregnant or breastfeeding, or may become pregnant or plan to breastfeed
- have severe kidney or liver problems
- are taking simvastatin or pravastatin

What is the most important safety information I should know about NEXLETOL?

NEXLETOL can cause side effects that can become serious, including:

- high levels of uric acid in the blood, which can lead to gout, a painful joint condition.

Call your doctor if you have any of the following symptoms:

- severe foot pain, especially in the toe joint
- warm joints
- swelling
- tender joints
- joint redness
- tendon problems: Stop taking NEXLETOL immediately and get medical help right away if you get any of the following signs or symptoms of tendon rupture:
 - hear or feel a snap or pop in a tendon area
 - bruising right after an injury in a tendon area
 - unable to move the affected area or put weight on the affected area

What are the common side effects of NEXLETOL?

Tell your doctor right away about any side effect you experience. NEXLETOL can cause side effects, including:

- symptoms of the common cold or flu
- muscle spasms
- back pain
- pain in the legs or arms
- stomach pain

Your doctor may check your blood for side effects that you might not be able to feel.

These are not all the possible side effects of NEXLETOL. For more information, ask your doctor or pharmacist.

What form does NEXLETOL come in?

NEXLETOL is a tablet that you take by mouth. It contains 180 mg of bempedoic acid.

Can I take NEXLETOL with other medicines?

For lowering cholesterol, NEXLETOL is often taken with other medicines, including a group of medicines called statins.

For two statins, simvastatin and pravastatin, your doctor may change the dose if you also take NEXLETOL.

Before taking NEXLETOL, tell your doctor if you take any of the following medicines:

- simvastatin
- pravastatin

The risk information provided here is not comprehensive. To learn more, talk about NEXLETOL with your doctor or pharmacist.

The FDA-approved product labelling can be found at [NEXLETOL.com](https://www.fda.gov/medwatch) or 1-833-377-7633.

You may report side effects to the FDA at 1-800-FDA-1088, or at www.fda.gov/medwatch.

You may also report side effects to Esperion at 1-833-377-7633.

NEXLETOL is available by prescription only.

Please see full [Prescribing Information](#).

ESPERION®

 **NEXLETOL®**
(bempedoic acid) tablets