

NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets

**Recommended by
the American Heart Association**



Can't take a statin? Help take control of your Lipid Lurkers (LDL-C) with NEXLIZET or NEXLETOL.

What are NEXLIZET and NEXLETOL?

- Along with diet and exercise:
 - NEXLETOL is used, with other cholesterol-lowering medicines, or alone when use with other cholesterol-lowering medicines is not possible, to reduce low-density lipoprotein (LDL) or bad cholesterol in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high blood cholesterol called heterozygous familial hypercholesterolemia.
 - NEXLIZET is used to reduce LDL in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high cholesterol called heterozygous familial hypercholesterolemia.
- Bempedoic acid, a component of NEXLIZET and NEXLETOL, is used to lower the risk of major adverse cardiovascular events, such as death from cardiovascular disease, heart attack, stroke, heart stent or bypass surgery in adults at increased risk for these events who are unable to take recommended statin treatment or are not taking a statin.

**Please see full Important Safety Information
on pages 8 and 9.**



How do I know if I'm statin intolerant?

If taking a statin caused side effects that led your doctor to lower your dose or you have stopped treatment, you may have statin intolerance. Talk to your doctor about nonstatin options to help lower your cholesterol.

You may not feel high cholesterol, but your arteries do.

Over time, Lipid Lurkers can build up and may cause a heart attack.



Who are the Lipid Lurkers?

They are your bad cholesterol (LDL-C) that often show no symptoms, so they pose a dangerous threat.

High cholesterol isn't just about what you eat. Most cholesterol is made by your body. While diet, exercise, and statins can help lower cholesterol, statins are not an option for everyone. If statin intolerance got in the way of your treatment, your bad cholesterol (AKA Lipid Lurkers), may still be putting you at risk.

IMPORTANT SAFETY INFORMATION

Do not take NEXLIZET if you are allergic to ezetimibe or bempedoic acid, or any ingredients in NEXLIZET or NEXLETOL. If you have any of the following signs and symptoms of a serious allergic reaction, call your healthcare provider or go to the nearest emergency room:

- swelling of your face, lips, mouth, or tongue
- wheezing
- severe itching
- fast heartbeat or pounding in your chest
- trouble breathing
- skin rashes, redness, or swelling
- dizziness or fainting

Did you know?

Up to 30% of patients are statin intolerant.

Without appropriate treatment, patients remain at a higher risk for a serious heart event, such as a heart attack.



If you're wondering what to do if you can't tolerate statins, NEXLIZET or NEXLETOL could be a statin alternative for you.

They work differently and are not active in the skeletal muscle.

NEXLETOL contains bempedoic acid.

NEXLIZET contains bempedoic acid and ezetimibe.

The CLEAR Outcomes trial studied the use of bempedoic acid exclusively in patients who can't take recommended statin therapy...

THE ONLY ORAL NONSTATIN STUDY for statin intolerant patients at high risk for a serious heart event.

Nearly 14,000 patients were studied to see if bempedoic acid could help reduce the risk of cardiovascular death, heart attack, stroke, heart stent, or bypass procedure.

Please see full Important Safety Information on pages 8 and 9.

NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets

How do I help take control of my Lipid Lurkers (LDL-C) when I can't take a statin?

NEXLIZET and NEXLETOL are the only oral nonstatins FDA approved to lower bad cholesterol and reduce the risk of the first or next serious heart event in patients with statin intolerance:

Serious heart events include:

- cardiovascular death
- heart attack
- stroke
- heart stent or bypass surgery



What are experts saying about using NEXLIZET or NEXLETOL as a statin alternative to lower bad cholesterol (LDL-C)?

According to the 2026 American College of Cardiology/American Heart Association (ACC/AHA) Guideline, people who cannot tolerate statins or need additional help lowering bad cholesterol (LDL-C) may benefit from proven nonstatin treatments, such as bempedoic acid.

IMPORTANT SAFETY INFORMATION

Before taking NEXLIZET or NEXLETOL, tell your healthcare provider if you:

- have or had gout
- have or had tendon problems
- are pregnant or breastfeeding or may become pregnant or plan to breastfeed. NEXLIZET or NEXLETOL may pass into your breastmilk. You and your healthcare provider should decide if you will take NEXLIZET or NEXLETOL or breastfeed.
- have moderate or severe liver problems

NEXLIZET and NEXLETOL are proven to help lower bad cholesterol along with a healthy diet and exercise.

NEXLIZET lowered bad cholesterol by an average of

38%



In a study, NEXLIZET lowered bad cholesterol by 36% compared to a sugar pill, which increased bad cholesterol by 2%.

NEXLETOL lowered bad cholesterol by an average of

~18%



In 2 studies, NEXLETOL lowered bad cholesterol by 15% in one trial and 17% in the second trial compared to a sugar pill, which increased bad cholesterol by 2%.



By taking NEXLIZET or NEXLETOL, patients should see the **maximum lowering of bad cholesterol by Week 4.**

Please see full Important Safety Information on pages 8 and 9.


NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets

NEXLIZET and NEXLETOL work differently than statins.

NEXLIZET and NEXLETOL are not statins. They interrupt your body's cholesterol-making process, lowering levels of LDL-C (AKA Lipid Lurkers).

NEXLETOL contains bempedoic acid.
NEXLIZET contains bempedoic acid and ezetimibe.



Is it easy to take NEXLIZET or NEXLETOL?

NEXLIZET or NEXLETOL can be used in combination with other cholesterol-lowering medications or alone.



NEXLIZET or NEXLETOL may be right for you.

Are you taking a statin but your Lipid Lurkers are not fully under control?

Have you stopped taking statins or are not able to take recommended statin therapy?

Talk to your healthcare provider to see if NEXLIZET or NEXLETOL is an option for you.

How do I take NEXLIZET or NEXLETOL?



Used with or without statins



One tablet, once a day



With or without food

Take 1 NEXLIZET or NEXLETOL tablet by mouth each day. Swallow the tablet whole.

If you miss a dose take it as soon as you remember. If you do not remember until it is time for your next dose, skip the dose you missed and continue with your regular schedule.

Do not take 2 doses at the same time.

Please see full Important Safety Information on pages 8 and 9.

NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements especially if you take or plan to take simvastatin, pravastatin, cyclosporine, fibrates, or cholestyramine.

Indication and Important Safety Information



What are NEXLIZET and NEXLETOL?

- Along with diet and exercise:
 - NEXLETOL is used, with other cholesterol-lowering medicines, or alone when use with other cholesterol-lowering medicines is not possible, to reduce low-density lipoprotein (LDL) or bad cholesterol in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high blood cholesterol called heterozygous familial hypercholesterolemia.
 - NEXLIZET is used to reduce LDL in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high cholesterol called heterozygous familial hypercholesterolemia.
- Bempedoic acid, a component of NEXLIZET and NEXLETOL, is used to lower the risk of major adverse cardiovascular events, such as death from cardiovascular disease, heart attack, stroke, heart stent or bypass surgery in adults at increased risk for these events who are unable to take recommended statin treatment or are not taking a statin.

Do not take NEXLIZET if you are allergic to ezetimibe or bempedoic acid, or any ingredients in NEXLIZET or NEXLETOL. If you have any of the following signs and symptoms of a serious allergic reaction, call your healthcare provider or go to the nearest emergency room:

- swelling of your face, lips, mouth, or tongue
- wheezing
- severe itching
- fast heartbeat or pounding in your chest
- trouble breathing
- skin rashes, redness, or swelling
- dizziness or fainting

Before taking NEXLIZET or NEXLETOL, tell your healthcare provider if you:

- have or had gout
- have or had tendon problems
- are pregnant or breastfeeding or may become pregnant or plan to breastfeed. NEXLIZET or NEXLETOL may pass into your breastmilk. You and your healthcare provider should decide if you will take NEXLIZET or NEXLETOL or breastfeed.
- have moderate or severe liver problems



Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements especially if you take or plan to take simvastatin, pravastatin, cyclosporine, fibrates, or cholestyramine.

NEXLIZET and NEXLETOL can cause serious side effects, including:

- **increased levels of uric acid in the blood**, which can lead to gout. Call your doctor if you have any of the following symptoms: severe foot pain, especially in the toe joint, warm joints, swelling, tender joints or joint redness.
- **tendon rupture or injury**. The risk of getting tendon problems while you take NEXLIZET or NEXLETOL is increased (1.2% vs 0.9% in placebo in Trial 1; 0.5% vs 0% in placebo in Trial 2). Stop taking NEXLIZET or NEXLETOL immediately and get medical help if you experience signs of tendon rupture such as snap or pop in tendon area, bruising after an injury in a tendon area, or unable to move or put weight on the affected area.

The most common side effects observed in trials of:

- **NEXLETOL includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, kidney problems, and gallstones.
- **NEXLIZET includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, diarrhea, joint pain, swelling of sinuses, fatigue, kidney problems, and gallstones.

These are not all the possible side effects of NEXLIZET and NEXLETOL. Ask your healthcare provider or pharmacist for more information.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see full Prescribing Information, including Patient Product Information, for NEXLIZET and NEXLETOL in pocket.

NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets

What are the side effects of NEXLIZET and NEXLETOL?

The combined safety of NEXLIZET and NEXLETOL has been demonstrated across multiple clinical trials.

The most common side effects observed in trials of:

- **NEXLETOL includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, kidney problems, and gallstones.
- **NEXLIZET includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, diarrhea, joint pain, swelling of sinuses, fatigue, kidney problems, and gallstones.

These are not all the possible side effects of NEXLIZET and NEXLETOL.

For more information, ask your healthcare provider or pharmacist.



What if I need help getting NEXLIZET or NEXLETOL?

If you have any questions or need help along the way, **NEXSTEP** Patient Support is here for you.

NEXSTEP

If your doctor prescribes NEXLIZET or NEXLETOL, the **NEXSTEP** Patient Support Program can:

- ✓ Answer treatment FAQs
- ✓ Help you understand why you may need cholesterol treatment
- ✓ Offer a Co-Pay Card to help with the cost of your prescription if you have commercial insurance



Scan to see if you're eligible for a Co-Pay Card, or find out at [NEXstep.com](https://www.nexstep.com)



Already have a prescription? Scan to enroll in **NEXSTEP** or visit [NEXLIZET.com/NEXSTEP](https://www.nexstep.com)

Please see full Important Safety Information on pages 8 and 9.

NEXLIZET[®]
(bempedoic acid/ezetimibe)
180mg/10mg tablets

NEXLETOL[®]
(bempedoic acid)
180mg tablets



Can't take a statin? Make NEXLIZET or NEXLETOL happen!

NEXLIZET and NEXLETOL are FDA approved to reduce the risk of serious heart events and lower bad cholesterol in patients with statin intolerance:

Serious heart events include:

- cardiovascular death
- heart attack
- stroke
- heart stent or bypass surgery

What are NEXLIZET and NEXLETOL?

- Along with diet and exercise:
 - NEXLETOL is used, with other cholesterol-lowering medicines, or alone when use with other cholesterol-lowering medicines is not possible, to reduce low-density lipoprotein (LDL) or bad cholesterol in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high blood cholesterol called heterozygous familial hypercholesterolemia.
 - NEXLIZET is used to reduce LDL in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high cholesterol called heterozygous familial hypercholesterolemia.
- Bempedoic acid, a component of NEXLIZET and NEXLETOL, is used to lower the risk of major adverse cardiovascular events, such as death from cardiovascular disease, heart attack, stroke, heart stent or bypass surgery in adults at increased risk for these events who are unable to take recommended statin treatment or are not taking a statin.



When statins alone aren't enough or you can't take them, consider NEXLIZET or NEXLETOL.

Please see full Important Safety Information on pages 8 and 9.

NEXLIZET[®]
(bempedoic acid/ezetimibe) tablets

NEXLETOL[®]
(bempedoic acid) tablets

ESPERION[™]

All trademarks and trade names are the property of their respective owners.

© 2026 Esperion Therapeutics, Inc. All rights reserved.
US-NXNZ-2600014