

TIPS FOR TALKING TO YOUR DOCTOR ABOUT

*If your bad cholesterol still isn't low enough,
you may be at risk for a serious heart event.[†]*

NEXLIZET and NEXLETOL are the only oral nonstatins* FDA approved to lower bad cholesterol and to reduce the risk of the first or next serious heart event[†] in patients with statin intolerance.

*Bempedoic acid

[†]Serious Heart Event = cardiovascular death, heart attack, stroke, heart stent or bypass surgery

To support you during your conversation with your doctor, print this form, complete it, and bring it to your next appointment.

My experience with lowering bad cholesterol

Fill in the below to help you and your doctor make a plan!

I have been diagnosed with high cholesterol:

☐ Yes ☐ No

My experience with statins is:

- ☐ I am currently taking a statin
- ☐ I have never taken a statin but would consider one, if recommended
- ☐ I stopped taking statins because of side effects or concerns of side effects
- ☐ I have never used a statin and do not want to take one

To lower my bad cholesterol, I have tried (check all that apply):

- ☐ A healthy diet
- ☐ Exercise
- ☐ Statins or other cholesterol-lowering medications
- ☐ Other: _____

I have had a heart attack, stroke, or heart procedure, such as stent placement or bypass surgery

☐ Yes (add details below)

☐ No
☐ Unsure

My current bad cholesterol (LDL-C) number is: _____ mg/dL
(If you don't know, ask your doctor)

My doctor recommends my bad cholesterol (LDL-C) number should be: _____ mg/dL
(If you don't know, ask your doctor)

I have been trying to lower my bad cholesterol for:

- ☐ 6 months–1 year ☐ 3 years–5 years
- ☐ 1 year–3 years ☐ 5+ years

Questions about my cholesterol

If you want to ask your doctor any of these questions, select them below.

- ☐ What is the difference between good cholesterol and bad cholesterol?
- ☐ Why is it important for me to know my number?
- ☐ What are my treatment options?

Questions to help you learn more about NEXLIZET or NEXLETOL

You can ask your doctor these questions to see if NEXLIZET or NEXLETOL may be right for you.

- How can NEXLIZET or NEXLETOL help lower my bad cholesterol (LDL-C)?
- How can NEXLIZET or NEXLETOL help reduce my risk of heart attack and heart procedures, like stent placement or bypass surgery?
- Why/how are NEXLIZET and NEXLETOL different from statins?
- What medicines are in NEXLIZET and NEXLETOL?
- What do I need to know about the safety of NEXLIZET or NEXLETOL?
- How do I take NEXLIZET or NEXLETOL?
- How often and for how long do I need to take NEXLIZET or NEXLETOL?
- Do NEXLIZET and NEXLETOL need to be taken with food?

IMPORTANT SAFETY INFORMATION

Do not take NEXLIZET if you are allergic to ezetimibe or bempedoic acid, or any ingredients in NEXLIZET or NEXLETOL. If you have any of the following signs and symptoms of a serious allergic reaction, call your healthcare provider or go to the nearest emergency room:

- swelling of your face, lips, mouth, or tongue
- fast heartbeat or pounding in your chest
- skin rashes, redness, or swelling
- wheezing
- trouble breathing
- dizziness or fainting
- severe itching

Please see additional Important Safety Information on the following page.

Additional questions and notes

It may also be helpful to bring a list of your current medications, recent lab results, and notes on any side effects you may be experiencing.

IMPORTANT SAFETY INFORMATION (continued)

Before taking NEXLIZET or NEXLETOL, tell your healthcare provider if you:

- have or had gout
- have or had tendon problems
- are pregnant or breastfeeding or may become pregnant or plan to breastfeed. NEXLIZET or NEXLETOL may pass into your breastmilk. You and your healthcare provider should decide if you will take NEXLIZET or NEXLETOL or breastfeed.
- have moderate or severe liver problems

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements especially if you take or plan to take simvastatin, pravastatin, cyclosporine, fibrates, or cholestyramine.

NEXLIZET and NEXLETOL can cause serious side effects, including:

- **increased levels of uric acid in the blood**, which can lead to gout. Call your doctor if you have any of the following symptoms: severe foot pain, especially in the toe joint, warm joints, swelling, tender joints or joint redness.
- **tendon rupture or injury**. The risk of getting tendon problems while you take NEXLIZET or NEXLETOL is increased (1.2% vs 0.9% in placebo in Trial 1; 0.5% vs 0% in placebo in Trial 2). Stop taking NEXLIZET or NEXLETOL immediately and get medical help if you experience signs of tendon rupture such as snap or pop in tendon area, bruising after an injury in a tendon area, or unable to move or put weight on the affected area.

The most common side effects observed in trials of:

- **NEXLETOL includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, kidney problems, and gallstones.
- **NEXLIZET includes** symptoms of the common cold or flu-like symptoms, muscle spasms, back pain, stomach pain, bronchitis, pain in shoulder, legs, or arms, anemia, increased liver enzymes, diarrhea, joint pain, swelling of sinuses, and fatigue.

These are not all the possible side effects of NEXLIZET and NEXLETOL. Ask your healthcare provider or pharmacist for more information.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What are NEXLIZET and NEXLETOL?

- Along with diet and exercise:
 - NEXLETOL is used, with other cholesterol-lowering medicines, or alone when use with other cholesterol-lowering medicines is not possible, to reduce low-density lipoprotein (LDL) or bad cholesterol in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high blood cholesterol called heterozygous familial hypercholesterolemia.
 - NEXLIZET is used to reduce LDL in adults with high blood cholesterol levels called hypercholesterolemia, including a type of high cholesterol called heterozygous familial hypercholesterolemia.
- Bempedoic acid, a component of NEXLIZET and NEXLETOL is used to lower the risk of major adverse cardiovascular events, such as death from cardiovascular disease, heart attack, stroke, stent placement or bypass surgery in adults at increased risk for these events who are unable to take recommended statin treatment or are not taking a statin

Please see full Prescribing Information, including Patient Product Information, for [NEXLIZET](#) and [NEXLETOL](#).

ESPERION

NEXLIZET[®]
(bempedoic acid/ezetimibe) tablets

NEXLETOL[®]
(bempedoic acid) tablets